

OPTIMIZING METABOLIC HEALTH AND NUTRITION IN THE TRANSITION COW

Sunita Lanje & Hrishikesh Tekade*

PhD Scholar, ICAR- Indian Veterinary Research Institute, Izatnagar, Bareilly Uttar Pradesh, India

Corresponding author's email: hrishikeshstekade2000@gmail.com

DOI: <https://doi.org/10.5281/zenodo.23448316>

Abstract

The transition period—encompassing the final three weeks before calving and the first three weeks after parturition—is universally recognized as the most metabolically and immunologically demanding phase of a dairy cow's lactation cycle. Driven by a sharp surge in nutrient requirements for milk synthesis and an accompanying depression in dry matter intake, cows undergo an obligate state of negative energy balance (NEB). Drawing from recent findings in ruminant science, this article highlights a paradigm shift in transition cow management: moving away from intensive prepartum energy loading ("steaming up") toward a structured, fundamentals-first approach. Key management strategies include securing an optimal body condition score (BCS) at calving, implementing moderate prepartum energy restriction (80% to 100% of calculated requirements), and providing targeted mineral supplementation, particularly for magnesium and calcium. Emphasizing practical field execution over complex precision protocols ensures reduced metabolic disease, improved milk production, and enhanced herd longevity.

Introduction

The transition from late gestation to early lactation represents a high-stress bottleneck in dairy production. Within 24 hours of parturition, the nutrient demands required to initiate milk synthesis multiply dramatically—reaching roughly 2.5 to 3 times the requirements of late pregnancy. Simultaneously, dry matter intake (DMI) typically drops by 30% to 50% right before calving.

This temporal mismatch creates an inevitable period of negative energy balance (NEB). If managed poorly, this metabolic strain triggers secondary disorders such as clinical and subclinical ketosis, hypocalcemia (milk fever), and fatty liver, contributing substantially to premature culling and economic loss.

For decades, standard extension advice recommended "steaming up" cows by feeding high levels of concentrates prior to calving to prime them for lactation. However, contemporary research—including recent reviews by Crookenden and Heiser (2026)—has overturned this paradigm. Evidence demonstrates that moderate prepartum energy restriction (feeding 80% to 100% of energy

requirements) actually "trains" the liver, improves metabolic health, and enhances immune competence compared to ad libitum or overfed regimes. This article explores how farmers and nutritionists can optimize metabolic health through practical, evidence-based management levers.

Physiological Realities of the Transition Phase

To manage transition cows effectively, producers must understand the coordinated homeorhetic adaptations that occur across multiple tissues weeks before and after calving:

- **Hepatic and Lipid Adaptations:** Gluconeogenic enzyme expression increases before and after calving to supply the massive glucose demand required for lactose synthesis. Simultaneously, nonesterified fatty acids (NEFA) mobilized from adipose tissue surge into the bloodstream, requiring efficient liver oxidation pathways.
- **Immunological Shifts:** Rather than a simple state of uniform immune "suppression," research indicates a coordinated shift from a

proinflammatory state prepartum to an anti-inflammatory profile immediately postpartum. Proper nutrition can significantly support the gene expression pathways responsible for neutrophil competence and oxidative stress responses.

Core Pillars of Transition Nutrition

A. Body Condition Score (BCS) as the Master Predictor

Achieving and maintaining target body condition scores is recognized as the single most robust predictor of transition success.

- The Targets: Mature cows should calve at a BCS of ~3.0 (on a 5-point scale, or 5.0 on a 10-point scale), while first-lactation heifers should target ~3.25 (5.5 on a 10-point scale).
- The Risks of Extremes: Cows calving at excessively high BCS (>3.25 / >5.5) experience severe lipolysis, heavy hepatic lipid accumulation, and a sharply elevated risk of ketosis. Conversely, underconditioned cows (<2.75 / <4.5) suffer from prolonged NEB, depressed milk yields, and impaired immunity.
- Management Timing: Management of BCS must be planned months in advance—typically during late lactation or the far-off dry period—rather than attempted right before calving.

B. Controlled Prepartum Energy Intake

Abandoning the old practice of heavy grain pre-calving has transformed transition diets.

- Restricting energy intake to 80% to 100% of calculated requirements during the final 3 to 4 weeks prepartum prevents exaggerated fat mobilization and reduces postpartum blood beta-hydroxybutyrate (BHB) concentrations.
- This controlled intake encourages hepatic gluconeogenic enzyme adaptation, effectively preparing the liver to process NEFA safely once milking begins.

C. Targeted Mineral Management (Magnesium and Calcium)

Mineral imbalances remain a primary gateway to metabolic collapse around parturition.

- Magnesium: Daily magnesium supplementation pre- and post-calving is vital to support smooth muscle function and prevent grass staggers or secondary deficiencies caused by high soil and pasture potassium levels.
- Calcium: Subclinical hypocalcemia is widespread around calving. Strategic calcium support during the colostrum period helps bridge the physiological gap when calcium demand for milk synthesis peaks abruptly. Rather than blanket-treating arbitrary laboratory cut-points, clinical intervention should focus on practical dietary support and targeted therapy for animals showing clear signs of weakness.

Emerging Horizons: Immunometabolism and Precision Tools

Modern science increasingly views the transition period through the lens of immunometabolism—acknowledging that metabolic stress directly dictates immune cell function.

- Inflammatory Regulation: Nutrients like omega-3 polyunsaturated fatty acids and antioxidant micronutrients (Vitamin E, selenium, zinc, copper) assist in downregulating harmful systemic inflammation and promoting tissue repair.
- Precision Monitoring: Wearable sensors tracking rumination time, feeding behavior, and 3-dimensional camera systems monitoring real-time body condition changes offer exciting pathways to flag at-risk cows days before clinical disease manifests. However, these technologies must be seamlessly integrated with practical farm labor realities and simple operational protocols

Conclusion

Optimizing transition cow performance does not require complex, high-input precision interventions. Instead, sustainable success relies on a fundamentals-first hierarchy centered around securing target body condition scores

well before calving, practicing controlled management with these core physiological prepartum energy feeding at 80% to 100% of requirements, and maintaining consistent, targeted magnesium and calcium supplementation. By aligning day-to-day farm realities, producers can drastically lower metabolic disease incidence, protect animal welfare, and secure long-term dairy profitability.

References

- Crookenden, M. A., A. Heiser. (2026). Transition cow nutrition for grazing herds: Practical priorities for health and performance. *JDS Communications* 7:34–40.
- Roche, J. R., A. W. Bell, T. R. Overton, J. J. Loor. (2013). Nutritional management of the transition cow in the 21st century—A paradigm shift in thinking. *Anim. Prod. Sci.* 53:1000–1023.
- Roberts, K. I., S. McDougall. (2019). Risk factors for subclinical hypocalcaemia, and associations between subclinical hypocalcaemia and reproductive performance, in pasture-based dairy herds in New Zealand. *N. Z. Vet. J.* 67:12–19.
- Horst, E. A., S. K. Kvidera, L. H. Baumgard. (2021). Invited review: The influence of immune activation on transition cow health and performance—A critical evaluation of traditional dogmas. *J. Dairy Sci.* 104:8380–8410.

Cite this article:

Sunita Lanje & Hrishikesh Tekade. (2026). Optimizing metabolic health and nutrition in the transition cow. *Vet Farm Frontier*, 03(09), 83–85. <https://doi.org/10.5281/zenodo.23448316>

